arXiv:cond-mat/0011118v1 [cond-mat.dis-nn] 7 Nov 2000

Dynamical Scaling Properties of Electrons in Quantum Systems with Multifractal Eigenstates

Jianxin Zhong,[1,2,3] Zhenyu Zhang,[2,1] Michael Schreiber[4], E. Ward Plummer,[1,2] and Qian Niu[2,5]
[1] *Department of Physics, University of Tennessee, Knoxville, Tennessee 37996*
[2] *Solid State Division, Oak Ridge National Laboratory, Oak Ridge, Tennessee 37831*
[3] *Department of Physics, Xiangtan University, Hunan 411105, P.R. China*
[4] *Institut für Physik, Technische Universität, D-09107 Chemnitz, Germany*
[5] *Department of Physics, University of Texas at Austin, Austin, Texas 78712*

We study the intricate relationships between the dynamical scaling properties of electron wave packets and the multifractality of the eigenstates in quantum systems. Numerical simulations for the Harper model and the Fibonacci chain indicate that the root mean square displacement displays the scaling behavior $r(t) \sim t^{\beta}$ with $\beta = D_2^{\psi}$, where D_2^{ψ} is the correlation dimension of the multifractal eigenstates. The equality can be generalized to d-dimensional systems as $\beta = D_2^{\psi}/d$, as long as the electron motion is ballistic in the effective D_2^{ψ}-dimensional space. This equality should be replaced by $\beta < D_2^{\psi}/d$ if the motion is non-ballistic, as supported by all known results.

PACS numbers: 71.23.Ft, 71.30.+h, 72.15.Rn 05.45.+b,

Spatial extension of eigenstates influences electron dynamics and transport properties of solids. It is well known that extended eigenstates in periodic lattices result in ballistic motion of electrons while localized eigenstates in strongly disordered systems lead to zero mobility. As a consequence, one expects metallic phase and insulating phase, respectively. In between these two extremes stand the critical eigenstates, i.e., the multifractal eigenstates, which are neither extended nor exponentially localized. Multifractal eigenstates exist in various quantum model systems, such as the Harper model of two-dimensional (2D) electrons in the presence of a perpendicular magnetic field [1], tight-binding models of quasicrystals [2–5], and disordered systems at the metal-insulator transition (MIT) [6–13]. In recent years, a great deal of effort has been devoted to the analysis of the multifractal nature of the eigenstates. Effort has also been made to establish intrinsic connections between the multifractality of the eigenstates and the transport properties in such systems. As an example, the unusual resistivity increase caused by decreasing temprature or improving structural quality of quasicrystals has been attributed to the multifractal nature of the electronic states [2]. Despite all these efforts, a fundamental level of understanding on precisely how the multifractality of the eigenstates affects the electron dynamics is still lacking.

The characteristics of electron dynamics is reflected by the time evolution of the corresponding wave packet, whose spreading can be described by the root mean square displacement $r(t)$. For ballistic spreading, $r(t) \sim t$, while in the absence of diffusion, $r(t) \sim t^0$ for large t. Numerical simulations [14] for the Harper model and the Fibonacci quasiperiodic chain [1–4] showed that $r(t) \sim t^{\beta}$ with $0 < \beta < 1$. Such a scaling behavior has also been found in other systems with multifractal eigenstates [10,12,15,16]. On the other hand, the energy spectrum of the multifractal eigenstates can be measured by the multifractal dimensions, D_q^{μ} [17]. The second moment, D_2^{μ}, has been shown to determine the decay of the average probability, $C(t)$, of a wave packet remaining at its initial position: $C(t) \sim t^{-D_2^{\mu}}$ [18,19]. It is naturally tempting to link the spectral multifractal dimensions, D_q^{μ}, with the dynamical scaling exponent, β [20–26]. Indeed, earlier such efforts have led to a variety of results, including $\beta \leq D_1^{\mu}$ [20], $\beta = D_{-1}^{\mu}$ [23,24], and $\beta \geq D_2^{\mu}/D_2^{\psi}$ [25], where D_2^{ψ} is the correlation dimension of the eigenstates. We note that all these results were derived with the assumption that the energy spectra are singularly continuous. Scaling behavior of $r(t)$ may also exist in higher dimensional systems whose spectra are not singularly continuous [15,16,19], but it would be even more difficult to derive a unique relation between β and D_q^{μ}.

In this Letter, we use a new approach to understand the scaling behavior of electron dynamics. Instead of focusing on the *energy* spectra, we study how the multifractality of the eigenstates in *real* space affects the time spreading of the wave packets. Our study leads to several fundamentally important findings. First, for the 1D Harper model and the Fibonacci chain, our numerical simulations show that β is directly given by the correlation dimension of the multifractal eigenstates: $\beta = D_2^{\psi}$. Second, by interpreting D_2^{ψ} as the effective dimension of the multifractal space for electron motion, we generalize the above relation to d-dimensional systems as $\beta = D_2^{\psi}/d$. This equality holds as long as the spreading of wave packets in the D_2^{ψ}-space is ballistic in nature, as demonstrated by applying to exactly solvable 2D and 3D quasiperiodic systems. Furthermore, when the spreading in D_2^{ψ}-space is non-ballistic, we have $\beta < D_2^{\psi}/d$, which is in complete agreement with all the known results for higher dimensional ($d > 1$) disordered systems [9–13].

We start with a 1D lattice of size $L = Na$, where N is the total number of sites and a the lattice constant. The Hamiltonian has the tight-binding form $H = \sum_n \epsilon_n |n >< n| + \sum_{m,n} t_{m,n} |m >< n|$, where ϵ_n is the site energy, $t_{m,n}$ the nearest-neighbor hopping elements. Eigen-energies and eigenstates are denoted as E_m and $\psi_n(E_m)$, respectively. The multifractal analysis of an eigenstate is based on the standard box-counting procedure: divide the system into a number of boxes of linear size wL $(0 < w < 1)$, and then calculate the box probability in the kth box, $p_k = \sum_{n \in k} |\psi_n(E_m)|^2$. A spatial correlation function $R(w, E)$ is defined as $R(w, E) = \sum_k p_k^2$. In general, $R(w, E) \sim w^{D_2^\psi(E)}$, where $D_2^\psi(E)$ is called the correlation dimension of the eigenstate. Extended eigenstates and localized eigenstates are characterized by $D_2^\psi(E) = 1$ and $D_2^\psi(E) = 0$, respectively. For multifractal eigenstates, $0 < D_2^\psi(E) < 1$. Considering the fact that eigenstates contributing to the time spreading of a wave packet may display different multifractal behaviors, we introduce a spectral-averaged spatial correlation function, $R(w) = \frac{1}{N}\sum_m R(w, E_m)$, and define a spectral-averaged correlation dimension D_2^ψ if $R(w) \sim w^{D_2^\psi}$ holds. The root mean square displacement for a wave packet initially located at site n_0 is given by $r(t, n_0) = \sqrt{\sum_n |n - n_0|^2 |\phi_n(t)|^2}$, where the wave function $\phi_n(t)$ can be obtained by $\phi_n(t) = \sum_m \psi_{n_0}^*(E_m)\psi_n(E_m)\exp(iE_m t)$. For inhomogeneous systems, $r(t, n_0)$ varies with n_0. In this case, one averages $r(t, n_0)$ over different initial wave packets in order to obtain the dynamics for the whole system. The scaling behavior of the system-averaged root mean square displacement is defined as $r(t) \sim t^\beta$, with $0 < \beta < 1$.



FIG. 1. Multifractal properties of eigenstates for the Harper model of system size $N = 3000$. The solid line shows $R(w) \sim w^{D_2^\psi}$ with $D_2^\psi = 0.5$. The dotted and the dashed lines show $R(w, E) \sim w^{D_2^\psi(E)}$ for energies E_e and E_c with $D_2^\psi(E_e) = 0.38$ and $D_2^\psi(E_c) = 0.64$, respectively.

Hamiltonian parameters in the Harper model [1] are $t_{m,n} = 1$ and $\epsilon_n = 2\cos(2\pi\sigma n)$, where $\sigma = (\sqrt{5} - 1)/2$ is the golden mean. We find that the correlation function $R(w, E)$ for various eigenstates displays a scaling behavior $R(w, E) \sim w^{D_2^\psi(E)}$, where $0 < D_2^\psi(E) < 1$ and $D_2^\psi(E)$ is different for different eigenstate. As examples, we show $R(w, E)$ for a system of $N = 3000$ in Fig. 1 at energies $E_\mathrm{e} = 2.59751453292$ and $E_\mathrm{c} = 0.00000066284$, which respectively correspond to the upper edge and the center of the energy spectrum. The corresponding scaling exponents are $D_2^\psi(E_\mathrm{e}) = 0.38$ and $D_2^\psi(E_\mathrm{c}) = 0.64$. We also find that the spectral-averaged (over the whole spectrum of $-2.6 < E < 2.6$) correlation function $R(w)$ displays a scaling behavior $R(w) \sim w^{D_2^\psi}$ with $D_2^\psi = 0.5$ as shown in Fig. 1. In Fig. 2, $R(w)$ is compared with $r(t)$ obtained by averaging over 500 initial wave packets. We find $r(t) \sim t^\beta$ with $\beta = 0.5$, in agreement with a previous study [24]. We therefore have $\beta = D_2^\psi = 0.5$.



FIG. 2. Scaling behaviors of $r(t) \sim t^\beta$ (thick lines) and $R(w) \sim w^{D_2^\psi}$ (thin lines) for the whole spectrum (solid lines), the center sub-spectrum (dashed lines), and the edge sub-spectrum (long dashed lines) in the Harper model with $\beta = D_2^\psi = 0.5$. System size is $N = 3000$.

We can also study how eigenstates in different spectral regions affect the spreading of wave packets by considering $\phi_n(t) = \sum_{E_m \in \Omega} \psi_{n_0}^*(E_m)\psi_n(E_m)\exp(iE_m t)$, where Ω denotes the spectral region. The corresponding $r(t)$ is compared with $R(w)$ obtained by averaging the eigenstates within Ω. Both the spectrum and the eigenstates of the Harper model have self-similar trifurcating structures [1–4]. Therefore, we expect that the spreading of the wave packets contributed by eigenstates in different sub-spectra follow the same scaling law. In Fig. 2, we show $R(w)$ and $r(t)$ for the center sub-spectrum $-0.2 < E < 0.2$ and the edge sub-spectrum $1.8 < E < 2.6$. It is clear that $R(w)$ for different sub-spectra is the same because of the same multifractal behavior of the eigenstates in different sub-spectra. Furthermore, $r(t)$ for different sub-spectra in Fig. 2 displays the same scaling law with $\beta = D_2^{\psi} = 0.5$, as expected. This indicates that the multifractality of the eigenstates indeed plays a crucial role in the spreading of the wave packets. We note that the downward shift of $r(t)$ shown in Fig. 2 for the sub-spectra is due to the decrease in the number of eigenstates contributing to the spreading of the wave packets.

We find the same relation $\beta = D_2^{\psi}$ to hold for the Fibonacci quasiperiodic chain [4]. In this model, $t_{i,j} = 1$ and ϵ_i takes $-u$ and u arranged in a Fibonacci sequence. The averaged correlation function has the scaling behavior $R(w) \sim w^{D_2^{\psi}}$ with $0 < D_2^{\psi} < 1$ depending on u. The system-averaged root mean square displacement $r(t)$ used to be compared with $R(w)$ was obtained by an average of 500 wave packets. As examples, we show $R(w)$ and $r(t)$ for the whole spectrum in Fig. 3 around $u = 1.65$: $\beta = D_2^{\psi} = 0.62$ for $u = 1$; $\beta = D_2^{\psi} = 0.5$ for $U = 1.65$; and $\beta = D_2^{\psi} = 0.43$ for $u = 2$. These three cases cover the super-diffusive, diffusive, and sub-diffusive regimes of motion in real space.



FIG. 3. Scaling behaviors of $r(t) \sim t^{\beta}$ (dashed lines) and $R(w) \sim w^{D_2^{\psi}}$ (solid lines) in the Fibonacci chain with site energies $u = 1, 1.65$, and 2. The corresponding exponents are $\beta = D_2^{\psi} = 0.62, 0.5$, and 0.43, respectively. System size is $N = 2584$.

The relation $\beta = D_2^{\psi}$ observed in different models can be generalized to higher dimensional systems with multifractal eigenstates. From its definition, $R(w)$ is in fact the average probability for an electron in a box of linear size wL. Because $R(w) \sim w^{D_2^{\psi}}$, the effective number of boxes of the system is $N' = (\frac{1}{w})^{D_2^{\psi}}$, instead of $(1/w)^d$, in order to ensure the normalization $N' \times R(w) \sim 1$. From the standard definition of the volume dimension D_v, i.e., *the number of the measured boxes*$\times(wL)^{D_v} =$ *constant*, we have $D_v = D_2^{\psi}$. Thus D_2^{ψ} can be viewed as the effective volume dimension for electron motion in the multifractal space (hereafter called the D_2^{ψ}-space). If the motion (i.e., the spreading of the wave packet) is ballistic in D_2^{ψ}-space, the volume, N', visited by the electron is $N' \sim t^{D_2^{\psi}}$. Because the width of the wave packet $r(t)$ is still measured in the d-dimensional real space, we have $r(t) \sim t^{D_2^{\psi}/d}$, which immediately leads to the result $\beta = D_2^{\psi}/d$.

As tests of the above generalized relation, we consider the exactly solvable models in higher dimensional Fibonacci lattices [15,19] and the Labyrinth lattices [27]. We find that the equality $\beta = D_2^{\psi}/d$ holds for all those models. As an example, here we only show the case for 3D Fibonacci lattice, which is constructed in a cubic lattice with constant hopping elements $t_{i,j} = 1$ and separable site energy $\epsilon_i = \epsilon_{ix} + \epsilon_{iy} + \epsilon_{iz}$, where $\epsilon_{ix}, \epsilon_{iy}$, and ϵ_{iz} are the 1D Fibonacci site energies. The energy spectrum is given by $E = E_x + E_y + E_z$ and the corresponding eigenstates are $\psi(E, x, y, z) = \psi_x(E_x) \times \psi_y(E_y) \times \psi_z(E_z)$, where E_j and $\psi_j(E_j)$ $(j = x, y, z)$ are the energy spectra and the eigenstates for the 1D Fibonacci chain, respectively. As a consequence, these eigenstates are multifractal [15,19], and their spreading dynamics is non-ballistic in real space: $r(t) \sim t^{\beta}$ [15], where $0 < \beta < 1$ takes the same value as that for the 1D Fibonacci chain. We note that all the relationships between the energy spectra and the dynamics [20–25] cannot give correct prediction for the spreading of wave packets in these higher dimensional quasiperiodic systems. In our approach, for the higher dimensional Fibonacci lattices, it is easy to show that $R(w)$ has the scaling behavior with $D_2^{\psi} = dD_2^{\psi_x}$, where $D_2^{\psi_x}$ is the multifractal dimension of the eigenstates in the 1D Fibonacci chain. Therefore, we have $\beta = D_2^{\psi}/d$.

Electron motion in the effective D_2^{ψ}-space can also be non-ballistic. In such cases, the above equality should be replaced by $\beta < D_2^{\psi}/d$, because no motion should be faster than ballistic. One special example is the diffusive motion ($\beta = 0.5$) in the metallic regime of 3D disordered systems with extended eigenstates ($D_2^{\psi} = 3$), where $\beta < D_2^{\psi}/3$. As shown below, this inequality is also satisfied by many other known examples involving non-ballistic motion in the D_2^{ψ}-space. At the metal-insulator transition of 3D disordered systems (in the absence of magnetic field), the eigenstates are found to be multifractal with different values of D_2^{ψ}: $D_2^{\psi}/3 = 0.57 \pm 0.1$ [6], $D_2^{\psi}/3 = 0.48 - 0.6$ [8], $D_2^{\psi}/3 = 0.57 \pm 0.07$ [9], and $D_2^{\psi}/3 = 0.5 \pm 0.07$ [10]. For such systems in a magnetic field, we have $D_2^{\psi}/3 = 0.57 \pm 0.07$ [10], $D_2^{\psi}/3 \approx 0.5$ [11], and $D_2^{\psi}/3 = 0.43 \pm 0.03$ [12]. On the other hand, the corresponding electron dynamics was found to have the same scaling exponent $\beta = 0.33$ [10], showing that $\beta < D_2^{\psi}/d$ for all these cases (with or without a magnetic field). Existing numerical results for 2D disordered systems satisfy this inequality as well. For the 2D disordered systems in a strong magetic field, eigenstates at the centers of the Landau bands are multifractal with $D_2^\psi/2 = 0.76 \pm 0.03$ and the corresponding spreading has a scaling behavior with $\beta = 0.5$ [12]. At the MIT of the 2D disordered systems with spin-orbit interactions, one finds $D_2^\psi/2 = 0.84 \pm 0.03$ and $\beta = 0.5$ [13]. It is evident that $\beta < D_2^\psi/d$.

In conclusion, we have shown that there exists an intrinsic relationship between the dynamic spreading of electron wave packets and the multifractality of the eigenstates in quantum systems. For the 1D Harper model and the Fibonacci chain, we have $\beta = D_2^\psi$, where β is the scaling exponent of the root mean square displacement, and D_2^ψ the correlation dimension of the multifractal eigenstates. We have further provided a conceptually new interpretation of D_2^ψ as the effective dimension for electron motion in the multifractal space. Following this interpretation, the above equality can be generalized to d-dimensional systems as $\beta = D_2^\psi/d$, as long as the electron motion is ballistic in the D_2^ψ-space. We have demostrated that such an equality applies to exactly solvable 2D and 3D quasiperiodic systems. When the motion in the D_2^ψ-space is non-ballistic due to disorder, the equality is replaced by $\beta < D_2^\psi/d$, which is in complete agreement with all the known results for higher dimensional ($d > 1$) disordered systems.

We thank T. Geisel, R. Ketzmerick, J. Bellissard, R.A. Römmer, U. Grimm, X.C. Xie and G. Canright for helpful discussions. This research was supported by the US National Science Foundation under Grant No. DMR-9702938, by Oak Ridge National Laboratory, managed by Lockheed Martin Energy Research Corp. for the Department of Energy under contract No. DE-AC05-96OR22464, and in part by the National Natural Science Foundation of China.

---

[1] P.G. Harper, Proc. Phys. Soc. London A **68**, 874 (1955); D.R. Hofstadter, Phys. Rev. B **14**, 2239 (1976); S. Aubry and G. Andre, Ann. Israel Phys. Soc. **3**, 133.

[2] D. Mayou, in *Lectures on Quasicrystals*, edited by F. Hippert and D. Gratias (Les Editions de Physique, Les Ulis, 1994), pp. 417-462; C. Berger, *ibid*, pp.463-504; C. Sire, *ibid*, pp.505-536.

[3] T. Fujiwara and H. Tsunetsugu, in *Quasicrystals: The State of the Art*, edited by D.P. Di Vincenzo and P.J. Steinhardt (World Scientific, Singapore, 1991).

[4] M. Kohmoto, L.P. Kadanoff, and C. Tang, Phys. Rev. Lett. **50**, 1870 (1983); S. Ostlund, R. Pandit, D. Rand, H.J. Schellnhuber, and E.D. Siggia, Phys. Rev. Lett. **50**, 1873 (1983); M. Kohmoto, B. Sutherland, and C. Tang, Phys. Rev. B **35**, 1020 (1987); Q. Niu and F. Nori, Phys. Rev. Lett. **57**, 2057 (1986).

[5] J.X. Zhong, U. Grimm, R.A. Römer, and M. Schreiber, Phys. Rev. Lett. **80**, 3996 (1998); J.X. Zhong and T. Geisel, Phys. Rev. E, in press (1999).

[6] C.M. Soukoulis and E.N. Economou, Phys. Rev. Lett. **52**, 565 (1984).

[7] F. Wegner, in *Localization and Metal-Insulator Transitions*, edited by H. Fritzsche and D. Adler, (Plenum Press, New York 1985), p. 337.

[8] M. Schreiber and H. Grussbach, Phys. Rev. Lett. **67**, 607 (1991).

[9] T. Brandes, B. Huckestein, and L. Schweitzer, Ann. Physik **5**, 633 (1996).

[10] T. Ohtsuki and T. Kawarabayashi, J. Phys. Soc. Jpn. **66**, 314 (1996).

[11] T. Terao, Phys. Rev. B **56**, 975 (1997).

[12] B. Huckestein and R. Klesse, cond-mat/9805038, (1998).

[13] L. Schweitzer, J. Phys. Conden. Matter **7**, L281 (1995); T. Kawarabayashi and T. Ohtsuki, Phys. Rev. B **53**, 6975 (1996).

[14] H. Hiramoto and S. Abe, J. Phys. Soc. Jpn. **57**, 230 (1998); *ibid* 1365 (1988).

[15] K. Ueda and H. Tsunetsugu, Phys. Rev. Lett. **58**, 1272 (1987).

[16] B. Passaro, C. Sire, and V.G. Benza, Phys. Rev. B **46**, 13751 (1992).

[17] T.G. Halsey, M.H. Jensen, L.P. Kadanoff, I. Procaccia and B.I. Shraiman, Phys. Rev. A **33**, 1141 (1986).

[18] R. Ketzmerick, G. Petschel, and T. Geisel, Phys. Rev. Lett. **69**, 695 (1992).

[19] J. X. Zhong and R. Mosseri, J. Phys. Condens. Matter **7**, 8383 (1995); J.X. Zhong, J. Bellissard, and R. Mosseri, J. Phys. Condens. Matter **7**, 3507 (1995).

[20] I. Guarneri, Europhys. Lett. **10**, 95 (1989); Europhys. Lett. **21**, (1993).

[21] T. Geisel, R. Ketzmerick, and G. Petschel, Phys. Rev. Lett. **66**, 1651 (1991).

[22] S. N. Evangelou and D. E. Katsanos, J. Phys. A **26**, L1243 (1993); M. Wilkinson and J. Austin, Phys. Rev. B **50**, 1420 (1994).

[23] G. Mantica, Physica (Amsterdam) **103D**, 576 (1997).

[24] F. Piechon, Phys. Rev. Lett. **76**, 4372 (1996).

[25] R. Ketzmerick, K. Kruse, S. Kraut, and T. Geisel, Phys. Rev. Lett. **79**, 1959 (1997).

[26] H. Schulz-Baldes and J. Bellissard, Rev. Math. Phys. **10**, 1 (1998).

[27] C. Sire, Europhys. Lett. **10**, 483 (1990).